



MESTEROLONE

MESTEROLONE 25 mg

Presented as box with 2 Blisters containing 15 capsules per carton

DESCRIPTION

Mesterolone is used to treat potency disturbances, infertility, declining physical & mental alertness in the aging male. It is an oral androgen which does not aromatize into estrogen. In clinical situations is generally used to treat various types of sexual dysfunction, which often result from a low endogenous testosterone level. It can usually reverse problems of sexual disinterest and impotency and is sometimes used to increase the sperm count. The drug does not stimulate the body to produce testosterone, but is simply an oral androgen substitute that is used to compensate for a lack of the natural male androgen. It is also effective act as an anti- aromatase in the body, actually preventing or slowing the conversion of androgens into estrogens. Mesterolone is approved for the following uses:

Declining physical activity and mental alertness in middle-aged and old men.

Hypogonadism- potency disturbances due to androgen deficiency. Infertility.

Oligozoospermia- to increase sperm count, improve quality and sperm motility.

CONTRAINDICATIONS

In-patients with carcinoma of the prostate, androgen therapy of any kind, including the use of Mesterolone is contraindicated.

PRECAUTIONS

The administration of Mesterolone is recommended only for male patients.

Regular examinations of the prostate should be carried out prophylactically.

INDICATIONS AND USAGE

Declining physical activity and mental alertness in middle and old aged men.

Reduced efficiency, easy fatigability, lack of concentration, weak memory, disturbances of libido and potency, irritability, disturbances of sleep, depressive moods, and general vegetative complaints are often attributed to androgen-deficiency.

Potency disturbances:

Mesterolone overcomes potency disturbances due to androgen deficiency. It may also be of use as supplementary therapy in cases of diminished potency where androgen-deficiency is not the primary cause.

Hypogonadism.

Growth, development, and function of androgen-dependent target organs are stimulated by Mesterolone. It promotes development of secondary male sex characteristics in cases of prepuberal hypogonadism. Full clinical and laboratory investigations are necessary in all cases of young patients prior to commencement of treatment.

Mesterolone 25 may also be used as a substitution therapy in cases where a loss of gonadal function has occurred post-puberally.

Infertility.

Oligozoospermia and deficient Leydig -cell secretion may be the cause of infertility. With Mesterolone treatment, sperm count can be increased, the quality improved and , furthermore, a higher fructose concentration up to normal values can be achieved thus increasing the chances of procreation.

PRESENTATION

25mg capsules in blister packs of 15 capsules — 2 blisters per box (30 capsules).

STORAGE

Store below 25°C. Store in the original package.