



TURINABOL

TURINABOL USP 10 mg

Presented as blister with 90 capsules

COMPOSITION

1 capsule contains: Active substance: 10 mg
4-chlorodehydromethyltestosterone
Magnesium stearate, Kollidon CL, FD & C
Yellow, Ludipress

PHARMACOLOGICAL PROPERTIES

Turinabol (4-chloro-17 α -dehydro-17 α -methyltestosterone) is considered to be an oral anabolic steroid mainly used for androgynous reduction. For this reason, androgynous activity is claimed to be 6%, while anabolic one is 53%. Turinabol, together with other anabolic and androgynous steroids, penetrates into the cell membrane and cytoplasmic receptors. When interacting they reach the nucleus where Turinabol affects certain nuclear receptors. The active substance combines with the DNA and performs as a transcription factor that causes the synthesis of specific RNA and proteins. Turinabol has mainly an anabolic effect which is mostly noticeable in the striated muscles and bones, together with other tissues with the intensive area (bone marrow, mucous membrane, etc). Its effect primarily determines the rate of muscle growth and mass increase. An essential contribution to the realization of metabolic diseases is considered to be the anabolic effect: positive nitrogen balance (nitrogen retention).

PHARMACOKINETIC PROPERTIES

The metabolism of Turinabol in the body takes from 6 to 8 days

INDICATIONS

Protein synthesis disorder, Traumas, Burns, Postoperative Infections, kidney failure, Osteoporosis, Corticosteroid therapy, Aplastic anemia, Negative nitrogen balance, Muscular atrophy when the patients.

DOSAGE AND HOW TO TAKE IT

The therapy with anabolic steroids is complementary and we can't replacing the primary treatment. The treatment depends from the patients and the side effects. Adults: the steroid effect is varied. The dose for adults is 20-50 mg for a day (2-6 weeks)

CONTRAINDICATIONS

Oversensitivity to the drug, prostate cancer, adenoma, breast cancer, Nephrotic syndrome, atherosclerosis, kidney failure, pregnancy, Breastfeeding.

OVERDOSE

There have been no reports of acute overdose with the androgens.

DRUG INTERACTIONS

When indirect anticoagulants are used together with anabolic steroids, their concentration and effects are reduced, so the dose needs to be reduced. It is necessary to monitor the prothrombin time. Concomitant use of ACTH and corticosteroids in patients with edema may worsen but the likelihood is very small. The free fraction of thyroid hormone levels remains unchanged

PRESENTATION

10 mg capsules in blister packs of 15 capsules - 6 blisters per box (90 capsules).

STORAGE

Store below 25°C. Store in the original package.